

1. Ikeda M, Aleksic B, Yamada K, Iwayama Y, Matsuo K, Numata S, Watanabe Y, Ohmura T, Kaneko T, Fukuo Y, Okochi T, Toyota T, Hattori E, Schibata S, Tanaka H, Yoneda H, Arai H, Someya T, Ohmori T, Yoshikawa T, Ozaki N, Iwata N: Genetic evidence for association between NOTCH4 and schizophrenia supported by a GWAS follow-up study in a Japanese population. *Mol Psychiatry* 2012-in press
2. Koide T, Banno M, Aleksic B (correspondence), Yamashita S, Kikuchi T, Kohmura K, Adachi Y, Kawano N, Kushima I, Nakamura Y, Okada T, Ikeda M, Ohi K, Yasuda Y, Hashimoto R, Inada T, Ujiike H, Iidaka T, Suzuki M, Takeda M, Iwata N, Ozaki N. Common variants in MAGI2 gene are associated with increased risk for cognitive impairment in schizophrenic patients. *PLoS One*. 2012-in press.
3. Uno Y, Uchiyama T, Kurosawa M, Aleksic B, Ozaki N. The combined measles, mumps, and rubella vaccines and the total number of vaccines are not associated with development of autism spectrum disorder: the first case-control study in Asia. *Vaccine*. 2012-in press.
4. Kitazawa M, Ohnuma T, Takebayashi Y, Shibata N, Baba H, Ohi K, Yasuda Y, Nakamura Y, Aleksic B, Yoshimi A, Okochi T, Ikeda M, Naitoh H, Hashimoto R, Iwata N, Ozaki N, Takeda M, Arai H. No associations found between the genes situated at 6p22.1, HIST1H2BJ, PRSS16, and PGBD1 in Japanese patients diagnosed with schizophrenia. *Am J Med Genet B Neuropsychiatr Genet*. 2012-in press
5. Koide T, Aleksic B (correspondence), Kikuchi T, Banno M, Kohmura K, Adachi A, Kawano N, Iidaka T, and Ozaki N Evaluation of Factors Affecting Continuous Performance Test Identical Pairs Version Score of Schizophrenic Patients in a Japanese Clinical Sample. *Schizophrenia Research and Treatment*. 2012, Article ID 970131,
6. Aleksic B, Kushima I, Hashimoto R, Ohi K, Ikeda M, Yoshimi A, Nakamura Y, Ito Y, Okochi T, Fukuo Y, Yasuda Y, Fukumoto M, Yamamori H, Ujiike H, Suzuki M, Inada T, Takeda M, Kaibuchi K, Iwata N, Ozaki N. Analysis of the VAV3 as Candidate Gene for Schizophrenia: Evidences From Voxel-Based Morphometry and Mutation Screening. *Schizophr Bull*. 2012-in press
7. Tamaji A, Iwamoto K, Kawamura Y, Takahashi M, Ebe K, Kawano N, Kunimoto S, Aleksic B, Noda Y, Ozaki N. Differential effects of diazepam, tandospirone, and paroxetine on plasma brain-derived neurotrophic factor level under mental stress. *Hum Psychopharmacol*. 2012-in press
8. Banno M, Koide T, Aleksic B (correspondence), Yamada K, Kikuchi T, Kohmura K, Adachi Y, Kawano N, Kushima I, Ikeda M, Inada T, Yoshikawa T, Iwata N, Ozaki N. A case control association study and cognitive function analysis of neuropilin and toll-like 1 gene and schizophrenia in the Japanese population. *PLoS One*. 2011;6(12):e28929.
9. Kikuchi T, Iwamoto K, Sasada K, Aleksic B, Yoshida K, Ozaki N. Sexual dysfunction and hyperprolactinemia in Japanese schizophrenic patients taking antipsychotics. *Prog Neuropsychopharmacol Biol Psychiatry*. 2012 Apr 27;37(1):26-32.
10. Ishikawa N, Goto S, Murase S, Kanai A, Masuda T, Aleksic B (correspondence), Usui H, Ozaki N. Prospective study of maternal depressive symptomatology among Japanese women. *J Psychosom Res*. 2011 Oct;71(4):264-9.
11. Iijima Y, Aleksic B, Ozaki N. Necessity for ethical consideration of research in the aftermath of disaster. *Psychiatry Clin Neurosci*. 2011 Aug;65(5):535-6.
12. Takahashi N, Nielsen KS, Aleksic B (equal contribution), Petersen S, Ikeda M, Kushima I, Vacaresse N, Ujiike H, Iwata N, Dubreuil V, Mirza N, Sakurai T, Ozaki N, Buxbaum JD, Sap J. Loss of function studies in mice and genetic association link receptor protein tyrosine phosphatase α to schizophrenia. *Biol Psychiatry*. 2011 Oct 1;70(7):626-35.
13. Kikuchi T, Iwamoto K, Sasada K, Aleksic B, Yoshida K, Ozaki N. Reliability and validity of a new sexual function questionnaire (Nagoya Sexual Function Questionnaire) for schizophrenic patients taking antipsychotics. *Hum Psychopharmacol*. 2011-in press
14. Yoshimura T, Usui H, Takahashi N, Yoshimi A, Saito S, Aleksic B, Ujiike H, Inada T, Yamada M, Uchimura N, Iwata N, Sora I, Iyo M, Ozaki N. Association analysis of the GDNF gene with methamphetamine use disorder in a Japanese population. *Prog Neuropsychopharmacol Biol Psychiatry*. 2011 Jul 1;35(5):1268-72.
15. Kushima I, Nakamura Y, Aleksic B, Ikeda M, Ito Y, Shiino T, Okochi T, Fukuo Y, Ujiike H, Suzuki M, Inada T, Hashimoto R, Takeda M, Kaibuchi K, Iwata N, Ozaki N. Resequencing and Association Analysis of the KALRN and EPHB1 Genes And Their Contribution to Schizophrenia Susceptibility. *Schizophr Bull*. 2010-in press
16. Ikeda M, Aleksic B (equal contribution), Kinoshita Y, Okochi T, Kawashima K, Kushima I, Ito Y,

- Nakamura Y, Kishi T, Okumura T, Fukuo Y, Williams HJ, Hamshere ML, Ivanov D, Inada T, Suzuki M, Hashimoto R, Ujike H, Takeda M, Craddock N, Kaibuchi K, Owen MJ, Ozaki N, O'Donovan MC, Iwata N. Genome-wide association study of schizophrenia in a Japanese population. *Biol Psychiatry*. 2011 Mar 1;69(5):472-8.
17. Nakamura Y, Ito Y, **Aleksic B**, Kushima I, Yasui-Furukori N, Inada T, Ono Y, Ozaki N. Influence of HTR2A polymorphisms and parental rearing on personality traits in healthy Japanese subjects. *J Hum Genet*. 2010 Dec;55(12):838-41.
18. Yoshimi A, **Aleksic B (correspondence)**, Kawamura Y, Takahashi N, Yamada S, Usui H, Saito S, Ito Y, Iwata N, Inada T, Noda Y, Yamada K, Ozaki N. Gene-wide association study between the methylenetetrahydrofolate reductase gene (MTHFR) and schizophrenia in the Japanese population, with an updated meta-analysis on currently available data. *Schizophr Res*. 2010 Dec;124(1-3):216-22.
19. Ogasawara K, Nakamura Y, **Aleksic B**, Yoshida K, Ando K, Iwata N, Kayukawa Y, Ozaki N. Depression associated with alcohol intake and younger age in Japanese office workers: a case-control and a cohort study. *J Affect Disord*. 2011 Jan;128(1-2):33-40.
20. Koide T, **Aleksic B**, Ito Y, Usui H, Yoshimi A, Inada T, Suzuki M, Hashimoto R, Takeda M, Iwata N, Ozaki N. A two-stage case-control association study of the dihydropyrimidinase-like 2 gene (DPYSL2) with schizophrenia in Japanese subjects. *J Hum Genet*. 2010 Jul;55(7):469-72.
21. **Aleksic B (correspondence)**, Kushima I, Ito Y, Nakamura Y, Ujike H, Suzuki M, Inada T, Hashimoto R, Takeda M, Iwata N, Ozaki N. Genetic association study of KREMEN1 and DKK1 and schizophrenia in a Japanese population. *Schizophr Res*. 2010 May;118(1-3):113-7.
22. Kushima I, **Aleksic B**, Ito Y, Nakamura Y, Nakamura K, Mori N, Kikuchi M, Inada T, Kunugi H, Nanko S, Kato T, Yoshikawa T, Ujike H, Suzuki M, Iwata N, Ozaki N. Association study of ubiquitin-specific peptidase 46 (USP46) with bipolar disorder and schizophrenia in a Japanese population. *J Hum Genet*. 2010Mar;55(3):133-6.
23. Kushima I, **Aleksic B (equal contribution)**, Ikeda M, Yamanouchi Y, Kinoshita Y, Ito Y, Nakamura Y, Inada T, Iwata N, Ozaki N. Association study of bromodomain-containing 1 gene with schizophrenia in Japanese population. *Am J Med Genet B Neuropsychiatr Genet*. 2010 Apr 5;153B(3):786-91.
24. Ikeda M, **Aleksic B**, Kirov G, Kinoshita Y, Yamanouchi Y, Kitajima T, Kawashima K, Okochi T, Kishi T, Zaharieva I, Owen MJ, O'Donovan MC, Ozaki N, Iwata N. Copy number variation in schizophrenia in the Japanese population. *Biol Psychiatry*. 2010 Feb 1;67(3):283-6.
25. Kawashima K, Ikeda M, Kishi T, Kitajima T, Yamanouchi Y, Kinoshita Y, Okochi T, **Aleksic B**, Tomita M, Okada T, Kunugi H, Inada T, Ozaki N, Iwata N. BDNF is not associated with schizophrenia: data from a Japanese population study and meta-analysis. *Schizophr Res*. 2009 Jul;112(1-3):72-9.
26. Ji X, Takahashi N, **Aleksic B**, Ishihara R, Nagai T, Mouri A, Saito S, Maeno N, Inada T, Ozaki N. An association between serotonin receptor 3B gene (HTR3B) and treatment-resistant schizophrenia (TRS) in a Japanese population. *Nagoya J Med Sci*. 2008 Mar;70(1-2):11-7.
27. **Aleksic B (correspondence)**, Ikeda M, Ishihara R, Saito S, Inada T, Iwata N, Ozaki N. No association between the oligodendrocyte-related gene PLP1 and schizophrenia in the Japanese population. *J Hum Genet*. 2008;53(9):863-6.
28. Kinoshita Y, Ikeda M, Ujike H, Kitajima T, Yamanouchi Y, **Aleksic B**, Kishi T, Kawashima K, Ohkouchi T, Ozaki N, Inada T, Harano M, Komiyama T, Hori T, Yamada M, Sekine Y, Iyo M, Sora I, Iwata N. Association study of the calcineurin A gamma subunit gene (PPP3CC) and methamphetamine-use disorder in a Japanese population. *Ann N Y Acad Sci*. 2008 Oct;1139:57-62.
29. Ito Y, Nakamura Y, Takahashi N, Saito S, **Aleksic B**, Iwata N, Inada T, Ozaki N. A genetic association study of the FXRD domain containing ion transport regulator 6 (FXRD6) gene, encoding phosphohippolin, in susceptibility to schizophrenia in a Japanese population. *Neurosci Lett*. 2008 Jun 13;438(1):70-5.
30. Saito S, Takahashi N, Maeno N, Ito Y, **Aleksic B**, Usui H, Iidaka T, Inada T, Ozaki N. An association study of tachykinin receptor 3 gene with schizophrenia in the Japanese population. *Neuroreport*. 2008 Mar 5;19(4):471-3.
31. Ikeda M, Takahashi N, Saito S, **Aleksic B**, Watanabe Y, Nunokawa A, Yamanouchi Y, Kitajima T, Kinoshita Y, Kishi T, Kawashima K, Hashimoto R, Ujike H, Inada T, Someya T, Takeda M, Ozaki N, Iwata N. Failure to replicate the association between NRG1 and schizophrenia using Japanese large sample. *Schizophr Res*. 2008Apr;101(1-3):1-8.
32. Yoshimi A, Takahashi N, Saito S, Ito Y, **Aleksic B**, Usui H, Kawamura Y, Waki Y, Yoshikawa T, Kato T, Iwata N, Inada T, Noda Y, Ozaki N. Genetic analysis of the gene coding for DARPP-32 (PPP1R1B) in Japanese patients with schizophrenia or bipolar disorder. *Schizophr Res*. 2008 Mar;100(1-3):334-41.
33. Maeno N, Takahashi N, Saito S, Ji X, Ishihara R, Aoyama N, **Aleksic B**, Miura H, Ikeda M, Suzuki T, Kitajima T, Yamanouchi Y, Kinoshita Y, Iwata N, Inada T, Ozaki N. Association of SOX10 with schizophrenia in the Japanese population. *Psychiatr Genet*. 2007 Aug;17(4):227-31. PubMed PMID: 17621166.

34. Maeno N, Takahashi N, Saito S, Ji X, **Aleksic B**, Ishihara R, Yoshida K, Inada T, Iidaka T, Ozaki N. Association study between the transferrin gene and schizophrenia in the Japanese population. *Neuroreport*. 2007 Mar 26;18(5):517-20.
35. **Aleksic B (correspondence)**, Ishihara R, Takahashi N, Maeno N, Ji X, Saito S, Inada T, Ozaki N. Gap junction coding genes and schizophrenia: a genetic association study. *J HumGenet*. 2007;52(6):498-501.
- 36.