

2012年 スタッフ業績一覧

Okada T, Sato W, Kubota Y, Toichi M, and Murai T. Right hemispheric dominance and interhemispheric cooperation in gaze-triggered reflexive shift of attention. *Psychiatry Clin Neurosci* 2012; 66 (2): 97-104

Watanabe Y, Egawa J, Iijima Y, Nunokawa A, Kaneko N, Shibuya M, Arinami T, Ujike H, Inada T, Iwata N, Tochigi M, Kunugi H, Itokawa M, Ozaki N, and Hashimoto R. Someya T. A two-stage case-control association study between the tryptophan hydroxylase 2 (TPH2) gene and schizophrenia in a Japanese population. *Schizophr Res* 2012; 137 (1-3): 264-6

Matsunaga S, Ikeda M, Kishi T, Fukuo Y, Aleksic B, Yoshimura R, Okochi T, Yamanouchi Y, Kinoshita Y, Kawashima K, Umene-Nakano W, Inada T, Kunugi H, Kato T, Yoshikawa T, Ujike H, Nakamura J, Ozaki N, Kitajima T, and Iwata N. An evaluation of polymorphisms in casein kinase 1 delta and epsilon genes in major psychiatric disorders. *Neurosci Lett* 2012; 529 (1): 66-9

Nakazaki C, Noda A, Koike Y, Yamada S, Murohara T, and Ozaki N. Association of insomnia and short sleep duration with atherosclerosis risk in the elderly. *Am J Hypertens* 2012; 25 (11): 1149-55

Horiuchi Y, Iida S, Koga M, Ishiguro H, Iijima Y, Inada T, Watanabe Y, Someya T, Ujike H, Iwata N, Ozaki N, Kunugi H, Tochigi M, Itokawa M, Arai M, Niizato K, Iritani S, Kakita A, Takahashi H, Nawa H, and Arinami T. Association of SNPs linked to increased expression of SLC1A1 with schizophrenia. *Am J Med Genet B* 2012; *Neuropsychiatr Genet* 2012; 159B (1): 30-7

Adachi Y, Aleksic B, Nobata R, Suzuki T, Yoshida K, Ono Y, and Ozaki N. Combination use of Beck Depression Inventory and two-question case-finding instrument as a screening tool for depression in the workplace. *BMJ Open* 2012; 2(3): 2:e000596

Koide T, Banno M, Aleksic B, Yamashita S, Kikuchi T, Kohmura K, Adachi Y, Kawano N, Kushima I, Nakamura Y, Okada T, Ikeda M, Ohi K, Yasuda Y, Hashimoto R, Inada T, Ujike H, Iidaka T, Suzuki M, Takeda M, Iwata N, and Ozaki N. Common Variants in MAGI2 Gene Are Associated with Increased Risk for Cognitive Impairment in Schizophrenic Patients. *PLoS One* 2012; 7 (5): e36836

Tamaji A, Iwamoto K, Kawamura Y, Takahashi M, Ebe K, Kawano N, Kunimoto S, Aleksic B, Noda Y,

- and Ozaki N. Differential effects of diazepam, tansospirone, and paroxetine on plasma brain-derived neurotrophic factor level under mental stress. *Hum Psychopharmacol* 2012; 27 (3): 329-33
- Nagashima W, Kimura H, Ito M, Tokura T, Arao M, Aleksic B, Yoshida K, Kurita K, and Ozaki N. Effectiveness of duloxetine for the treatment of chronic nonorganic orofacial pain. *Clin Neuropharmacol* 2012; 35 (6): 273-7
- Torii Y, Iritani S, Sekiguchi H, Habuchi C, Hagikura M, Arai T, Ikeda K, Akiyama H, and Ozaki N. Effects of aging on the morphologies of Heschl's gyrus and the superior temporal gyrus in schizophrenia: A postmortem study. *Schizophr Res* 2012; 134 (2-3): 137-42
- Kawano N, Iwamoto K, Ebe K, Suzuki Y, Hasegawa J, Ukai K, Umegaki H, Iidaka T, and Ozaki N. Effects of mild cognitive impairment on driving performance in older drivers. *J Am Geriatr Soc* 2012; 60 (7): 1379-81
- Koide T, Aleksic B, Kikuchi T, Banno M, Kohmura K, Adachi Y, Kawano N, Iidaka T, and Ozaki N. Evaluation of factors affecting continuous performance test identical pairs version score of schizophrenic patients in a Japanese clinical sample. *Schizophr Res Treatment* 2012; 2012: 970131
- Ohi K, Hashimoto R, Yasuda Y, Fukumoto M, Yamamori H, Umeda-Yano S, Okada T, Kamino K, Morihara T, Iwase M, Kazui H, Numata S, Ikeda M, Ohnuma T, Iwata N, Ueno S, Ozaki N, Ohmori T, Arai H, and Takeda M. Functional genetic variation at the NRGN gene and schizophrenia: Evidence from a gene-based case-control study and gene expression analysis. *Am J Med Genet B Neuropsychiatr Genet* 2012; 159B (4): 405-13
- Kishi T, Ichinose H, Yoshimura R, Fukuo Y, Kitajima T, Inada T, Kunugi H, Kato T, Yoshikawa T, Ujike H, Musso G, M, Umene-Nakano W, Nakamura J, Ozaki N, and Iwata N. GTP cyclohydrolase 1 gene haplotypes as predictors of SSRI response in Japanese patients with major depressive disorder. *J Affect Disord* 2012; 142 (1-3): 315-22
- Kitazawa M, Ohnuma T, Takebayashi Y, Shibata N, Baba H, Ohi K, Yasuda Y, Nakamura Y, Aleksic B, Yoshimi A, Okochi T, Ikeda M, Naitoh H, Hashimoto R, Iwata N, Ozaki N, Takeda M, and Arai H. No associations found between the genes situated at 6p22.1, HIST1H2BJ, PRSS16, and PGBD1 in Japanese patients diagnosed with schizophrenia. *Am J Med Genet B Neuropsychiatr Genet* 2012; 159B (4): 456-64

- Kimura H, Yoshida K, Ito M, Tokura T, Nagashima W, Kurita K, and Ozaki N. Plasma levels of milnacipran and its effectiveness for the treatment of chronic pain in the orofacial region. *Hum Psychopharmacol* 2012; 27 (3): 322-8
- Furumura K, Koide T, Okada T, Murase S, Aleksic B, Hayakawa N, Shiino T, Nakamura Y, Tamaji A, Ishikawa N, Ohoka H, Usui H, Banno N, Morita T, Goto S, Kanai A, Masuda T, and Ozaki N. Prospective Study on the Association between Harm Avoidance and Postpartum Depressive State in a Maternal Cohort of Japanese Women. *PLoS One* 2012; 7 (4): e34725
- Tokura T, Kimura H, Yoshimi A, Ohashi M,S, Masuda M, Senzaki K, Yoshida K, Aleksic B, Noda Y, Yamada K, and Ozaki N. Reliability and Validity of the Japanese Version of BEMIB Modified for Patients With Bipolar Disorder: a Self-rating Scale for Medication Adherence. *Clinical Neuropsychopharmacology and Therapeutics* 2012; 3: 26-32
- Kushima I, Nakamura Y, Aleksic B, Ikeda M, Ito Y, Shiino T, Okochi T, Fukuo Y, Ujike H, Suzuki M, Inada T, Hashimoto R, Takeda M, Kaibuchi K, Iwata N, and Ozaki N. Resequencing and Association Analysis of the KALRN and EPHB1 Genes And Their Contribution to Schizophrenia Susceptibility. *Schizophr Bull* 2012; 38 (3): 552-60
- Kishi T, Fukuo Y, Okochi T, Kawashima K, Kitajima T, Inada T, Ozaki N, Musso G, M, Kane J, M, Correll C, U, and Iwata N. Serotonin 6 receptor gene and schizophrenia: case-control study and meta-analysis. *Hum Psychopharmacol* 2012; 27 (1): 63-9
- Kikuchi T, Iwamoto K, Sasada K, Aleksic B, Yoshida K, and Ozaki N. Sexual dysfunction and hyperprolactinemia in Japanese schizophrenic patients taking antipsychotics. *Prog Neuropsychopharmacol Biol Psychiatry* 2012; 37 (1): 26-32
- Kawano N, Iwamoto K, Ebe K, Aleksic B, Noda A, Umegaki H, Kuzuya M, Iidaka T, and Ozaki N. Slower adaptation to driving simulator and simulator sickness in older adults. *Aging Clin Exp Res* 2012; 24 (3): 285-9
- Uno Y, Uchiyama T, Kurosawa M, Aleksic B, and Ozaki N. The combined measles, mumps, and rubella vaccines and the total number of vaccines are not associated with development of autism spectrum disorder: The first case-control study in Asia. *Vaccine* 2012; 30 (28): 4292-8
- Hayakawa N, Koide T, Okada T, Murase S, Aleksic B, Furumura K, Shiino T, Nakamura Y, Tamaji A, Ishikawa N, Ohoka H, Usui H, Banno N, Morita T, Goto S, Kanai A, Masuda T, and Ozaki N.

The postpartum depressive state in relation to perceived rearing: a prospective cohort study.
PLoS One 2012; 7 (11): e50220

Hagikura M, Iwamoto K, Aleksic B, and Ozaki N. What is a rational antidepressant treatment for major depression in patients with Parkinson's disease? *Psychiatry Clin Neurosci* 2012; 66 (5): 463
Iidaka T, Miyakoshi M, Harada T, and Nakai T. White matter connectivity between superior temporal sulcus and amygdala is associated with autistic trait in healthy humans. *Neuroscience Letters* 2012; 510 (2): 154-8

Iidaka T, Harada T, Kawaguchi J, and Sadato N. Neuroanatomical substrates involved in true and false memories for face. *NeuroImage* 2012; 62 (1): 167-76

Iidaka T, Harada T, Eifuku S, Nakata R, and Sadato N. Distinct human face representations in the perirhinal cortex and fusiform gyrus. *Brain Research* 2012; 1452: 119-29

Banno M, Koide T, Aleksic B, Okada T, Kikuchi T, Kohmura K, Adachi Y, Kawano N, Iidaka T, and Ozaki N. Wisconsin Card Sorting Test scores and clinical and sociodemographic correlates in Schizophrenia: multiple logistic regression analysis. *BMJ Open* 2012; 2 (6): e001340

Suzuki M, Takahashi S, Matsushima E, Tsunoda M, Kurachi M, Okada T, Hayashi T, Ishii Y, Morita K, Maeda H, Katayama S, Otsuka T, Hirayasu Y, Sekine M, Okubo Y, Motoshita M, Ohta K, Uchiyama M, and Kojima T. Relationships between exploratory eye movement dysfunction and clinical symptoms in schizophrenia. *Psychiatry Clin Neurosci* 2012; 66 (3): 187-94