

2013年 スタッフ業績一覧

- Niwa M, Jaaro-Peled H, Tankou S, Seshadri S, Hikida T, Matsumoto Y, Cascella N. G, Kano S, Ozaki N, Nabeshima T, Sawa A. Adolescent stress-induced epigenetic control of dopaminergic neurons via glucocorticoids. *Science* 2013; 339 (6117): 335-9
- Takata A, Iwayama Y, Fukuo Y, Ikeda M, Okochi T, Maekawa M, Toyota T, Yamada K, Hattori E, Ohnishi T, Toyoshima M, Ujike H, Inada T, Kunugi H, Ozaki N, Nanko S, Nakamura K, Mori N, Kanba S, Iwata N, Kato T, Yoshikawa T. A Population-Specific Uncommon Variant in GRIN3A Associated with Schizophrenia. *Biol Psychiatry* 2013; 73 (6): 532-9
- Furukawa M, Tochigi M, Otowa T, Arinami T, Inada T, Ujike H, Watanabe Y, Iwata N, Itokawa M, Kunugi H, Hashimoto R, Ozaki N, Kakiuchi C, Kasai K, Sasaki T. An association analysis of the cardiomyopathy-associated 5 (CMYA5) gene with schizophrenia in a Japanese population. *Psychiatr Genet* 2013; 23 (4): 179-80
- Aleksic B, Kushima I, Hashimoto R, Ohi K, Ikeda M, Yoshimi A, Nakamura Y, Ito Y, Okochi T, Fukuo Y, Yasuda Y, Fukumoto M, Yamamori H, Ujike H, Suzuki M, Inada T, Takeda M, Kaibuchi K, Iwata N, Ozaki N. Analysis of the VAV3 as Candidate Gene for Schizophrenia: Evidences From Voxel-Based Morphometry and Mutation Screening. *Schizophr Bull* 2013; 39 (3): 720-8
- Kamijima K, Higuchi T, Ishigooka J, Ohmori T, Ozaki N, Kanba S, Kinoshita T, Koyama T, Group AS. Aripiprazole augmentation to antidepressant therapy in Japanese patients with major depressive disorder: A randomized, double-blind, placebo-controlled study (ADMIRE study). *J Affect Disord* 2013; 151 (3): 899-905
- Manchia M, Adli M, Akula N, Ardaur R, Aubry JM, Backlund L, Banzato CE, Baune BT, Bellivier F, Bengesser S, Biernacka JM, Brichant-Petitjean C, Bui E, Calkin CV, Cheng AT, Chillotti C, Cichon S, Clark S, Czerski PM, Dantas C, Zompo MD, Depaulo JR, Detera-Wadleigh SD, Etain B, Falkai P, Frisén L, Frye MA, Fullerton J, Gard S, Garnham J, Goes FS, Grof P, Gruber O, Hashimoto R, Hauser J, Heilbronner U, Hoban R, Hou L, Jamain S, Kahn JP, Kassem L, Kato T, Kelsoe JR, Kittel-Schneider S, Kliwicki S, Kuo PH, Kusumi I, Laje G, Lavebratt C, Leboyer M, Leckband SG, López Jaramillo CA, Maj M, Malafosse A, Martinsson L, Masui T, Mitchell PB, Mondimore F, Monteleone P, Nallet A, Neuner M, Novák

- T, O'Donovan C, Osby U, Ozaki N, Perlis RH, Pfennig A, Potash JB, Reich-Erkelenz D, Reif A, Reininghaus E, Richardson S, Rouleau GA, Rybakowski JK, Schalling M, Schofield PR, Schubert OK, Schweizer B, Seemüller F, Grigoriu-Serbanescu M, Severino G, Seymour LR, Slaney C, Smoller JW, Squassina A, Stamm T, Steele J, Stopkova P, Tighe SK, Tortorella A, Turecki G, Wray NR, Wright A, Zandi PP, Zilles D, Bauer M, Rietschel M, McMahon FJ, Schulze TG, Alda M. Assessment of Response to Lithium Maintenance Treatment in Bipolar Disorder: A Consortium on Lithium Genetics (ConLiGen) Report. *PLoS One* 2013; 8 (6): e65636
- Shiino T, Koide T, Kushima I, Ikeda M, Kunimoto S, Nakamura Y, Yoshimi A, Aleksic B, Banno M, Kikuchi T, Kohmura K, Adachi Y, Kawano N, Okada T, Inada T, Ujike H, Iidaka T, Suzuki M, Iwata N, Ozaki N. Common variants in bcl9 gene and schizophrenia in a japanese population: Association study, meta-analysis and cognitive function analysis. *J Med Biochem* 2013; 32 (4): 351-7
- Aleksic B, Kushima I, Ohye T, Ikeda M, Kunimoto S, Nakamura Y, Yoshimi A, Koide T, Iritani S, Kurahashi H, Iwata N, Ozaki N. Definition and refinement of the 7q36.3 duplication region associated with schizophrenia. *Sci Rep* 2013; 3: 2587
- Tanaka S, Syu A, Ishiguro H, Inada T, Horiuchi Y, Ishikawa M, Koga M, Noguchi E, Ozaki N, Someya T, Kakita A, Takahashi H, Nawa H, Arinami T. DPP6 as a candidate gene for neuroleptic-induced tardive dyskinesia. *Pharmacogenomics J* 2013; 13 (1): 27-34
- Ukai K, Kimura H, Arai M, Aleksic B, Yamauchi A, Ishihara R, Iritani S, Kurita K, Ozaki N. Effectiveness of low-dose milnacipran for a patient suffering from pain disorder with delusional disorder (somatic type) in the orofacial region. *Psychogeriatrics* 2013; 13 (2): 99-102
- Iwamoto K, Kawano N, Sasada K, Kohmura K, Yamamoto M, Ebe K, Noda Y, Ozaki N. Effects of low-dose mirtazapine on driving performance in healthy volunteers. *Hum Psychopharmacol* 2013; 28 (5): 523-8
- Sasada K, Iwamoto K, Kawano N, Kohmura K, Yamamoto M, Aleksic B, Ebe K, Noda Y, Ozaki N. Effects of repeated dosing with mirtazapine, trazodone, or placebo on driving performance and cognitive function in healthy volunteers. *Hum Psychopharmacol* 2013; 28 (3): 281-6
- Kohmura K, Iwamoto K, Aleksic B, Sasada K, Kawano N, Katayama H, Noda Y, Noda A, Iidaka T, Ozaki N. Effects of sedative antidepressants on prefrontal cortex activity during verbal

- fluency task in healthy subjects: a near-infrared spectroscopy study. *Psychopharmacology (Berl)* 2013; 226 (1): 75-81
- Ikeda M, Okahisa Y, Aleksic B, Won M, Kondo N, Naruse N, Aoyama-Uehara K, Sora I, Iyo M, Hashimoto R, Kawamura Y, Nishida N, Miyagawa T, Takeda M, Sasaki T, Tokunaga K, Ozaki N, Ujike H, Iwata N. Evidence for shared genetic risk between methamphetamine-induced psychosis and schizophrenia. *Neuropsychopharmacology* 2013; 38 (10): 1864-70
- Seo S, Takayama K, Uno K, Ohi K, Hashimoto R, Nishizawa D, Ikeda K, Ozaki N, Nabeshima T, Miyamoto Y, Nitta A. Functional Analysis of Deep Intronic SNP rs13438494 in Intron 24 of PCLO Gene. *PLoS One* 2013; 8 (10): e76960
- Esaki K, Kondo K, Hatano M, Saito T, Kishi T, Umene-Nakano W, Yoshimura R, Nakamura J, Ozaki N, Ikeda M, Iwata N. Further evidence of an association between a genetic variant in BMP7 and treatment response to SSRIs in major depressive disorder. *J Hum Genet* 2013; 58 (8): 568-9
- Ikeda M, Aleksic B, Yamada K, Iwayama-Shigeno Y, Matsuo K, Numata S, Watanabe Y, Ohnuma T, Kaneko T, Fukuo Y, Okochi T, Toyota T, Hattori E, Shimodera S, Itakura M, Nunokawa A, Shibata N, Tanaka H, Yoneda H, Arai H, Someya T, Ohmori T, Yoshikawa T, Ozaki N, Iwata N. Genetic evidence for association between NOTCH4 and schizophrenia supported by a GWAS follow-up study in a Japanese population. *Mol Psychiatry* 2013; 18 (6): 636-8
- Kondo K, Ikeda M, Kajio Y, Saito T, Iwayama Y, Aleksic B, Yamada K, Toyota T, Hattori E, Ujike H, Inada T, Kunugi H, Kato T, Yoshikawa T, Ozaki N, Iwata N. Genetic Variants on 3q21 and in the Sp8 Transcription Factor Gene (SP8) as Susceptibility Loci for Psychotic Disorders: A Genetic Association Study. *PLoS One* 2013; 8 (8): e70964
- Hashimoto R, Ikeda M, Ohi K, Yasuda Y, Yamamori H, Fukumoto M, Umeda-Yano S, Dickinson D, Aleksic B, Iwase M, Kazui H, Ozaki N, Weinberger DR, Iwata N, Takeda M. Genome-wide association study of cognitive decline in schizophrenia. *Am J Psychiatry* 2013; 170 (6): 683-4
- Shibata H, Yamamoto K, Sun Z, Oka A, Inoko H, Arinami T, Inada T, Ujike H, Itokawa M, Tochigi M, Watanabe Y, Someya T, Kunugi H, Suzuki T, Iwata N, Ozaki N, Fukumaki Y. Genome-wide association study of schizophrenia using microsatellite markers in the Japanese population. *Psychiatr Genet* 2013; 23 (3): 117-23

- Balan S, Yamada K, Iwayama Y, Toyota T, Ohnishi T, Maekawa M, Toyoshima M, Iwata Y, Suzuki K, Kikuchi M, Ujike H, Inada T, Kunugi H, Ozaki N, Iwata N, Nanko S, Kato T, Yoshikawa T. Lack of association of EGR2 variants with bipolar disorder in Japanese population. *Gene* 2013; 526 (2): 246-50
- Asano M, Esaki K, Wakamatsu A, Kitajima T, Narita T, Naitoh H, Ozaki N, Iwata N. Maternal overprotection score of the Parental Bonding Instrument predicts the outcome of cognitive behavior therapy by trainees for depression. *Psychiatry Clin Neurosci* 2013; 67 (5): 340-4
- Kawano N, Awata S, Ijuin M, Iwamoto K, Ozaki N. Necessity of normative data on the Japanese version of the Wechsler Memory Scale-Revised Logical Memory subtest for old-old people. *Geriatr Gerontol Int* 2013; 13 (3): 726-30
- Miyata S, Noda A, Iwamoto K, Kawano N, Okuda M, Ozaki N. Poor sleep quality impairs cognitive performance in older adults. *J Sleep Res* 2013; 22 (5): 535-41
- Watanabe Y, Iijima Y, Egawa J, Nunokawa A, Kaneko N, Arinami T, Ujike H, Inada T, Iwata N, Kunugi H, Itokawa M, Sasaki T, Ozaki N, Hashimoto R, Shibuya M, Igeta H, Someya T. Replication in a Japanese population that a MIR30E gene variation is associated with schizophrenia. *Schizophr Res* 2013; 150 (2-3): 596-7
- Ukai K, Okajima A, Yamauchi A, Sasaki E, Yamaguchi Y, Kimura H, Aleksic B, Ozaki N. Total palliative care for a patient with multiple cerebral infarctions that occurred repeatedly in association with gastric cancer (Trousseau's syndrome). *Palliat Support Care* 2013; 11 (2): 169-72
- Fujishiro H, Iseki E, Kasanuki K, Chiba Y, Ota K, Murayama N, Sato K. A follow up study of non-demented patients with primary visual cortical hypometabolism: prodromal dementia with Lewy bodies. *J Neurol Sci* 2013; 334 (1-2): 48-54
- Fujishiro H, Imamura AY, Lin WL, Uchikado H, Mark MH, Golbe LI, Markopoulou K, Wszolek ZK, Dickson DW. Diversity of pathological features other than Lewy bodies in familial Parkinson's disease due to SNCA mutations. *Am J Neurodegener Dis* 2013; 2 (4): 266-75
- Usami M, Okada T, Sasayama D, Iwadare Y, Watanabe K, Ushijima H, Kodaira M, Sugiyama N, Sawa T, Saito K. What time periods of the day are concerning for parents of children with attention deficit hyperactivity disorder? *PLoS One* 2013; 8 (11): e79806
- Fujishiro H, Kasanuki K, Nakamura S. Levodopa treatment and mood fluctuation in dementia with Lewy bodies: a case report. *Psychogeriatrics* 2013; 13 (4): 250-3

- Iritani S. What happens in the brain of schizophrenia patients?: an investigation from the viewpoint of neuropathology. *Nagoya J Med Sci* 2013; 75 (1-2): 11-28
- Fujishiro H, Iseki E, Nakamura S, Kasanuki K, Chiba Y, Ota K, Murayama N, Sato K. Dementia with Lewy bodies: early diagnostic challenges. *Psychogeriatrics* 2013; 13 (2): 128-38
- Kamio Y, Inada N, Moriwaki A, Kuroda M, Koyama T, Tsujii H, Kawakubo Y, Kuwabara H, Tsuchiya, KJ, Uno Y, Constantino JN. Quantitative autistic traits ascertained in a national survey of 22,529 Japanese schoolchildren. *Acta Psychiatrica Scandinavica* 2013; 128 (1): 45-53
- Kasanuki K, Iseki E, Nishida Y, Fujishiro H, Chiba Y, Sato K, Arai H. Effectiveness of ramelteon for visual hallucinations in dementia with Lewy bodies: four cases report. *J Clin Psychopharmacol* 2013; 33 (4): 581-3
- Chiba Y, Katsuse O, Fujishiro H, Kamada A, Saito T, Ikura T, Takahashi Y, Kunii M, Takeno M, Hirayasu Y. Lymphopenia helps early diagnosis of systemic lupus erythematosus for patients with psychosis as an initial symptom. *Psychosomatics* 2013; 3182: 136-9